

Garry K. Smith¹

¹Newcastle and Hunter Valley Speleological Society Inc.
P.O. Box 15, Broadmeadow, N.S.W. 2292, Australia. gksmith29@icloud.com

Abstract

The generic term ‘rootsicle’ has been used for several decades to describe plant roots in caves that are coated in a secondary mineral deposit. Rootsicles are found in shallow caves worldwide and take on a variety of forms and mineral coatings, but most commonly calcite. In caves, rootsicles can take on similar forms to stalactites, columns and stalagmites. There are at least three types of interactions that can take place between roots and secondary cave deposits, one of which results in a form that cannot be considered a rootsicle. There can be large variations in morphology and petrology between rootsicles growing at a cave entrance, to those in the twilight and dark zones. Influencing factors can include microclimate around developing rootsicles (temperature, humidity) and also light intensity. However, rootsicle-like forms are not restricted to caves. Above-ground plant roots can also become coated in secondary minerals and the resulting structure can look very similar to those formed underground. While above-ground forms are obviously not speleothems, it is not unreasonable to also describe them as rootsicles given similarities in their forms and in the multiplicity of influences on their development. A revised definition is therefore proposed to better capture the current knowledge surrounding the formation of rootsicles and associated structures.

Introduction

Rootsicle is the generic term embraced by speleologists for all forms of plant roots that have grown into a cave, been coated in a secondary mineral, usually calcium carbonate (CaCO_3) and thus effectively become fossilised (Hill and Forti 1997; Smith 1999; Hill 1999). More recently, the definition was expanded to include speleothems, produced by the coating or replacement of plant roots (Taboroši and others 2004). This process can result in formation of a column, stalactite or stalagmite cast in the shape of the roots, often having a twisted or contorted shape (Figure 1; Hill and Forti 1997; Taboroši 2004, 2006).

Rootsicles are less common than other forms of secondary deposits in caves. In general, only tree roots are able to grow deep enough to enter shallow caves (up to 30 m below the surface) and have a chance to become rootsicles over time. However, fig and boab tree roots have been observed at depths between 40 and 50 m in karst areas across Northern Australia, which are subject to seasonally humid climate (Gillieson 2004, Smith 2018).

Rootsicles should not be confused with ‘rhizoliths’ which are organosedimentary structures resulting from the preservation of roots of higher plants, or the remains thereof, in soils and sedimentary deposits rather than within caves (Klappa 1980).



Figure 1. Rootsicle embedded into flowstone in Cow Cave (MC-46), Mole Creek, Tasmania. Photo: Garry K. Smith

Creation of rootsicles

Tree roots grow into cave voids in search of water and nutrients, then become rootsicles if they are coated in CaCO_3 (or another mineral – see ‘Non-carbonate Rootsicles’, below) deposited

Rootsicles, roots & caves

from seepage water splashing onto or running down them. The roots may become a preferential area for the flow of seeping water and for calcium carbonate deposition (Shopov 2004). CaCO_3 is usually deposited as 'calcite' as this is a more stable polymorph, than aragonite and vaterite. Provided the supply of seepage water continues, the calcite layers may become so thick that the roots are completely encased and can't be seen (Figure 2).



Figure 2. A broken section of a rootsicle shows consecutive layers of calcite which have grown around a group of small roots. The roots had decomposed and their remnants washed out of the speleothem prior to natural breakage under its own weight. Photo: Garry K. Smith

If the calcite layer is not too thick and the roots continue to grow, the calcite may crack off the roots in strips and fall to the cave floor to form what can look like a pile of sticks. If the tree roots die, the calcite layers may continue to grow thicker, and although the roots will slowly decay they may still provide sufficient support to the developing speleothem until it is strong enough to stand-alone. The slow breakdown of the dead roots occurs because the fungi, mould and other organisms that decay the encapsulated roots are starved of oxygen. Eventually the rotting roots will decay to a consistency of papier-mâché, often jet black in colour (Smith 2010a). If sufficient water then enters these root canals, their contents can be washed out to leave just hollow tubes, with the internal shape resembling the original roots. Well after the roots have decayed the outside may continue to be coated in successive layers of calcite and the deposit preserves the roots approximate shape and original form.

When they form inside caves, they are commonly referred to by the informal term "rootsicles", although not every root-associated speleothem should be so termed. There are at least three different types of interactions that can take place between roots and secondary cave deposits, see Table 1. The resultant speleothems are distinct from each other and do not always fit the definition of rootsicle. Type 3, Table 1, is not a rootsicle because in this case the speleothem has formed first, then the root has grown through the solution path, cracks or porous voids in the speleothem (Taboroši and others 2004). Because the root has not been coated in a secondary deposit it can't be considered a rootsicle.

In their study of caves in southern Thailand Taboroši and others (2005) found a large variation in morphology and petrology between stalactites (including rootsicles) growing at a cave entrance and those in the twilight and dark zones. Microclimate (temperature, humidity) and light intensity were found to influence the morphology of stalactites and rootsicles. At the cave entrance largely biogenic stalactitic accretions of calcareous tufa were found growing around the drip-line (and even outside the cave) and there was a transition in their morphology to the dense coarsely crystalline stalactites (speleothems) in the cave's interior. Around the cave entrance, irregular, porous and easily recognizable tufa fabrics had formed around hanging plant roots, "resulting in tufaceous equivalents of cave rootsicles" (Figure 3). They noted that in tropical cave entrances, stalactites are soft and fragile in the most exposed locations, and more dense and solid in better-enclosed areas (Taboroši and others 2005).

In general, when a stalactite and stalagmite join, there are usually characteristic indications in the shape of the resulting column. This is often a step change in the column diameter (larger at the bottom or vice versa) and also a significant change in the external shape. However, when a tree root extends from ceiling to floor in a cave, and then becomes coated in a thick layer of calcite, the resulting rootsicle column is likely to be roughly uniform in diameter for its full length. Rootsicle columns may

Table 1 - Types of root-speleothem interactions in terms of basic temporal relationships between the plant roots and calcite deposits.




Temporal relationship	1	2	3
	Roots precede speleothems	Roots and speleothems develop concurrently	Speleothems precede roots
Sequence diagram			
Effects of roots	Constructional	Constructional	Destructional
Typical deposits	Rootsicles	Tufaceous stalactites	Stal. invaded by roots

Table 1. From Taboroši and others, 2004. Reproduced with permission.



Figure 3. Roots and rootsicles at the entrance of Pop Kan Mai Cave in Chong Phli village, Ao Nang subdistrict of Krabi, Thailand. Both stalactites are rootsicles, i.e. roots coated in a tufaceous calcite and aragonite deposit. The one on the left has more algae and moss growing on its surface. Photo: Danko Taboroši.

also consist of quite a few roots which have grown closely together from ceiling to floor. Thus, the shape may provide a clue to its origin as tree roots that created a path for seepage water to run down.

Breakdown, Damage and Repair

Rootsicles consisting of fine calcite coated roots that have rotted and fallen from the main roots, may form a mass resembling tangled spaghetti (Hill and Forti 1997). They may also look like a pile of calcified ‘fiddle sticks’ on the floor when relatively straight thin roots have rotted and the calcite structure has broken under its own weight.

Larger rootsicles that are accidentally damaged or naturally break under their own weight, typically result in a pile of many broken pieces (Figure 4). If broken due to natural causes rootsicles are typically left untouched, however if wilfully or accidentally broken attempts may be made to restore them. If attempts are made to repair humanly-caused damage to rootsicles it can prove very difficult because their centres are usually hollow after the



Figure 4. Broken rootsicle in Wildmans Cave (W-456), Wombeyan, NSW. Photo: Garry K. Smith.

Rootsicles, roots & caves

roots decay, thus the surface area at the break points is usually of insufficient extent to provide a strong bonding surface for adhesives. Cleaning out the rotten tree root canal for some distance into the broken segments and inserting threaded pins with water resistant adhesives up the central holes, can achieve limited success (Smith 2010a). Details of environmentally friendly adhesives are available elsewhere (Smith 2010a).

Occurrences of Rootsicles

Rootsicles can be found in many caves around Australia. For example, at Wombeyan (NSW) a 2.5 m calcified tree root has formed a column in Shawl Cave (W-12) (James and others 1982) and another 2 m long (now broken) rootsicle stalactite was formerly present in Wildmans Cave (W-456) (Smith 2009, 2010a). At the entrance to Wollondilly (W-144) and Creek (W-149) caves there are distorted, irregular stalactites that appear to be calcified tree roots (James and others 1982). Rootsicles in other NSW caves include Stove Pipe (WA-5) at Walli Caves, (Smith 2001); a 30 cm rootsicle in Gaden Caves (WE-2) at Wellington (Ian Eddison pers. comm. 2022); and a 3.5 m rootsicle forming a column in Death Trap Cave at Glenrock (GR-124) (Smith 2010b). In Rock-me Cave at Timor, NSW there are slender rootsicle stalactites ~5 m in length (Figure 5) (Baker 2007; Rutledge 2008).

Giants Cave (WI-21), south of Margaret River, WA contains several very impressive calcified bunches of tree roots hanging from the ceiling and referred to locally as arborites (rootsicles) (Smith 2003). Bastian (2014) states that these speleothems (arborites) are very common in caves developed in aeolian calcarenite along Australia's western seaboard. The roots in caves, north and south of Perth WA are mostly of the Tuart tree (*Eucalyptus gomphocephala*), which has an affinity for limestone.

Overseas, rootsicles up to 5 m long have been reported as being prevalent in caves of Mexico's Yucatan peninsula, where they may form the basis of most of the columns that form in these shallow dry caves (Bunnell 2021). They are also found in subsequently drowned caves, such as Dos Pisos Cenote (Two Levels Cenote) in the Yucatan region, which has large rootsicle columns in excess of 3 m high, at a depth of ~5.5 m below the present water surface (Figure 6). Images provided by cave



Figure 5. Andrew Baker next to a >5m long delicate rootsicle in Rock-Me Cave (TR-52) Timor, NSW. Photo: Garry K. Smith.



Figure 6. Submerged rootsicles in Dos Pisos Cenote (Two Levels Cenote) - Yucatan region. Photo: Neil Vincent.

diving photographer Neil Vincent, show well-formed stalactites, and other speleothems including rootsicles that were created during a period when the cave passages were dry. Many of the presently flooded caves and cenotes were dry during the Last Glacial Maximum at the peak of the Ice Age (about 22,000 years ago) when the sea level was 120

meters below its current level, postglacial sea level rise then causing back-flooding of many of caves (González-González and others 2008; Blanchon and Shaw 1995). The calcium saturated cave water has preserved the rootsicles for thousands of years.

There are some fine examples of calcite rootsicles in Caverne de La Vierge on the Mauritian island of Rodrigues (G. Middleton, pers. comm.) (Figure 7). The roof thickness above the cave is about 10 m and the passage where the main rootsicles occur is about 1.5 m high. The karst is developed in aeolian calcarenite very similar to that found on Australia's west coast.



Figure 7. Rootsicle column in Caverne de La Vierge, Rodrigues Island, Mauritius. Photo: G.J. Middleton.

Rootsicle Stalagmites

Tree roots can do strange things and have been observed growing upwards from the cave floor in search of moisture. For example, some roots in Monbulla Cave (L-5) near Naracoorte in SA (Figures 8 and 9) grow upwards from the earth floor and have been partly coated in calcite (Smith 2007). The roots are growing upward beneath a localised drip point in search of water. In both examples the lower halves of the central roots are completely encased in calcite, yet fine rootlets continue to grow out of the top and side of the calcite casing. They are considered rootsicle stalagmites under the current definition.



Figures 8 and 9. Upward growing rootsicles beneath drip points in Monbulla Cave (L-5) SA. Note AA battery for scale. Photos: Garry K. Smith.

In Deepdene Cave (AU-1) in WA, tree-root-entwined stalagmites were reported by De Waele (2014), whose report includes a photo depicting a stalagmite that appears to be at least 40 cm high with a considerable amount of partly calcite-encrusted rootlets covering most of it. From the photo it appears that the stalagmite has grown first, and the fine roots have grown up the outside and have been gradually encased in calcite. Again, this raises the question - Is it, or is it not a rootsicle?

Du Preez and others (2015) described what they called, 'hairy stalagmites', a new type of speleothem, which had only been observed in a couple of Botswana caves (Dimapo and Diviner's caves)

Rootsicles, roots & caves

which were previously sealed. The description of these (Namaqua fig tree) root stalagmites almost completely coated in calcite, fits the definition of rootsicles. Their stereo and electron microscopy study revealed the structures to consist of multiple intertwined tubes created when thin films of CaCO_3 are deposited around fine lateral roots. The hairy stalactites ranged in height from just a few centimetres to more than a metre, and their diameter varied between 3 and 5 cm.

Globally, rootsicle stalagmites (speleothems) appear to be quite rare, however the occurrence of roots growing upwards to create a structure resembling a stalagmite, appears to be more common than was thought by authors of early papers (Winkelhofer 1975; Lamont and Lange 1976). While not rootsicles, they have the potential to be transformed if seepage water contains minerals that may be deposited on them.

Winkelhofer (1975) was the first to describe root stalagmites from sandstone caves in Germany. These were described as conical and/or cylindrical dense networks of roots. Voids between the main roots in the stalagmite structures consisted mainly of living terminal roots often coated by symbiotic fungi, and were filled with sandy grains and organic matter. No calcite crust was deposited on the root stalagmites, due to the chemical composition of the host rock. The height of these structures was up to 60 cm.

Lamont and Lange (1976) described vertically growing fibrous root structures occurring in ten limestone caves near Yanhep National Park, Western Australia. They called these structures ‘Stalagmiform’ roots and determined that they were the roots of large trees (*Eucalyptus gomphocephala*) growing above the caves. The tallest structure was 12 cm and a few had calcite deposited from dripwater amongst the root structures.

Marais and others (1996), reported that in Pofaddergat Cave, Namibia, there are rootlets having a stalagmitic form (root stalagmites), which have preferentially grown immediately below dripping sites. “They form small stalagmites of densely packed thin roots, which are not more than a couple of centimetres in diameter and up to 20 cm high”. From the description there appears to be no calcite deposition on the roots.

Mlejnek (2010) identified that in late 2009 there were 78 localities with 245 root formations

documented (mostly stalagmites) in the Czech Republic. In the Poseidon sandstone cave system, root stalagmites grew up to 60 cm high and 10 cm or more in diameter (Mlejnek and others 2008). “In the rest of the world there are only 38 [root stalagmite] localities documented (Poland, Germany, Austria, Slovakia, Hungary, Sweden, Spain, South Africa, Australia) and in these localities 81 root stalagmites and stalactites were discovered” (Mlejnek 2010). By far the majority of these were found in sandstone talus caves. In 2009 only a handful of root stalagmites were reported in carbonate caves: one known location in the Czech Republic’s Moravian Karst, one cave in Austria, three caves in Hungary, and one cave in Australia (Mlejnek and others 2008, 2009; Mlejnek 2010).

Non-carbonate Rootsicles

Speleothem rootsicles have also been reported in Galeria da Queimada lava tube, on the Portuguese island of Terceira in the Azores archipelago (Daza Brunet and Bustillo 2014). The rootsicles have a mineralogical composition and developmental association with biomineralization induced by microbial activity in caves. Three types of rootsicle were defined, namely white hard and black spongy types, both composed of allophane, and a third hard, red coloured type composed of hydrous ferric oxy-hydroxide minerals (Daza Brunet and Bustillo 2014).

Speleothems such as flowstone, stalactites, stalagmites and coralline can occur in other non-carbonate caves. Rootsicles can be found in quartz sandstone overhangs and shallow caves of the Central Coast, Sydney and Blue Mountains regions of New South Wales. They can be created when calcium carbonate is leached from the overlying sandstone strata (Wray 1995, Smith 2015). Alternatively, they may be created from the deposition of iron oxides/hydroxides, or opal-A interlayered with chalcedony. They are physically robust and range in colour from white through to orange and almost black, depending on the chemical composition and proportion of included organic matter (Wray 1995, 1999). Figure 10 shows bright orange rootsicles coated in iron oxides/hydroxides and algae at the entrance to a small overhang cave in the Watagan Mountains, NSW.

“Related forms” (Hill and Forti 1997) are those deposits that resemble rootsicles, but are not speleothems in the strictest sense because they are not composed of true minerals but of mud or organic material



Figure 10. Rootsicles coated in iron oxides/hydroxides and algae, at the entrance to a small overhang cave in the Watagan Mountains. They are physically robust and range in colour from bright orange through to almost black. Photo: Garry K. Smith.

Roots Above and Below Ground

In the tropical and monsoonal areas of Australia, tree roots can grow to an enormous size. Such examples can be found in the Northern Territory, at Bullita caves, where boab and fig trees extend their roots down into the caves in search of water and nutrients (Figures 11 and 12). The wet and dry seasons mean that most of the ~550 mm average annual rainfall occurs in just 4 or 5 months (approx. between November and April) while little or no rain falls during the rest of the year. It is during the dry periods when the trees particularly need long roots to extract moisture from soil beneath the cave floor. When it rains the trees on the karst tend to catch droplets causing water to preferentially flow down their trunks, then follow the root systems into the cave. Despite this there is typically little calcite deposited on the tree roots because of the high flow rates when it rains. The calcium carried in solution is transported out of the cave and discharged at effluxes into surface perennial creeks flowing toward major river systems. In places so much calcite is carried out of the cave systems that tufa dams are created along surface stream beds, due to evaporation and degassing from the solution. Deposits similar to rootsicles can form on grasses, twigs and roots, mainly on the downstream side of tufa dams (Figure 13).



Figure 11. In Two Fishes Cave (BAA-11), Bullita, NT, large tree roots penetrate to great depths in search of water. Photo: Garry K. Smith.



Figure 12. Melissa Hadley next to tree roots that are flooded during the wet season and completely dry the rest of the year, in Neighbours Block (BAA-51) Bullita NT. Note the water-level mark on roots. Photo: Garry K. Smith.



Figure 13. Heiko Maurer and above-ground rootsicles at Calcite Waterfall, Gregory National Park, NT. Photo: Garry K. Smith.

In regions with less rainfall or with soils and porous rock which quickly drain water away, trees and other vegetation have adapted by sending roots down into caves where they hang from their ceilings like drapery. These fine rootlet masses hang in the cave's high humidity atmosphere, absorbing droplets which condense on them (Smith 2007). An example is in L-23 cave (incorrectly called Quarry Cave by Smith 2007) near Naracoorte, SA (Figures 14 and 15).



Figure 14. Jessica Bayles amongst roots absorbing condensation from a high humidity atmosphere in L-23 near Naracoorte, SA. Photo: Garry K. Smith.

Tree roots in caves play an important role in the subterranean food chain and can support a wide range of fauna, discussion of which is beyond the intended scope of this article.

From a safety aspect, speleologists should be aware that roots can be a source of elevated CO_2 in a cave. Plant root respiration can elevate CO_2 to dangerous concentrations in cave atmospheres where there is little air movement to disperse it. Also, microbial and fungal activity, breaking down dead roots, will reduce oxygen and increase CO_2 concentrations in caves.

Discussion

At what point does a plant root become a rootsicle and does it only apply to speleothems? These are the questions that the author grappled with when writing this article. Current definitions don't clearly define the percentage of tree root that must be coated in a secondary mineral deposit (e.g. calcite), to make it a rootsicle. Is it $>50\%$ coated, and how thick should the coating be? The root could be partly encased with a hard coating over 25% of its length and the rest might have just a visible thin white film, which may be powdery (e.g. Figure 16). What happens if a root is completely encased for its full length and is considered a rootsicle for many years and then a branching root breaks out of the calcite casing and continues to grow without a coating? Is there a point where the rootsicle ceases to be one, because the root has grown a certain percentage longer than the original rootsicle?

Defining this speleothem can be quite difficult as there are so many variables. Even setting an arbitrary percentage of secondary deposit coating and thickness on roots is prob-




Figure 15. Fine rootlets suspended from cave ceiling, capture condensation in L-23 cave near Naracoorte SA. Photo: Garry K. Smith.



Figure 16. The two groups of tree root either side of caver are about 50% coated in a layer of calcite and the remainder is a very thin powdery coating. Are they rootsicles? The roots on far left have no coating. Reto Zollinger in Dingo Cave, Bullita, NT. Photo: Garry K. Smith

lematic when it comes to accurate measurement. The easiest solution might be to leave the definition open to interpretation, by saying it becomes a rootsicle if part of the root is coated in a secondary mineral deposit.

If a rootsicle can only be a speleothem, there is also the dilemma of, can it be a rootsicle if outside a cave? A cave is typically defined in conventional lay dictionaries as being “a natural void in the ground, specifically a space large enough for a human to enter”, whereas in most speleological literature the definition also includes “must have a dark zone”. So, if a cave fits the description in speleological literature and has numerous rootsicles from the back of the cave (dark zone) right through to the drip line at the entrance (daylight), is there a point where they should not be considered a rootsicle? An overhang (large enough for a human to enter) under a waterfall can be considered a cave according to the lay dictionary definition, but not according to a speleological definition. If there are roots coated in a secondary mineral deposit under the drip line (Figure 10), are they considered to be in a cave or not? The answer appears to depend on how pedantic one is in interpreting and applying the definition of a cave.

Then there is the situation when a mixture of roots and vegetation is coated in a secondary deposit above ground (outside the cave environment) (Figure 13). Should these still be called ‘rootsicles’ even though they cannot be considered ‘speleothems’? The term “speleothem” as introduced by Moore (1952), is derived from the Greek words *spēlaion* ‘cave’ + *thēma* ‘deposit’, which specifically defines speleothems as secondary deposits formed in a cave.

Even in their acclaimed book *Cave Minerals of the World*, Hill and Forti (1997), leave room for interpretation between their descriptions of rootsicles on page 224 and their definition on page 363. Their description suggests that the root becomes covered with CaCO_3 so that it, “may be obscured or decayed, leaving only a calcium carbonate cast of the root”, while their definition says the root becomes calcified. Since the term rootsicle is a ‘generic term’ for roots covered in a secondary mineral deposit (typically calcite), it makes sense that similar occurrences above ground should be called the same thing, however above ground they can’t be considered a speleothem. This would also align with the generic terms ‘stalactite’ and ‘stalagmite’,

Rootsicles, roots & caves

which are often used to describe the shape of a secondary mineral deposit or structure in non-cave locations such as on cliff faces, and hence which are not necessarily speleothems even though speleothems are the most common form of stalactite.

Bastian (2014) argues that 'rootsicles' should be called 'arborites'. His reasoning is that, "speleothems generally have the suffix '-ite', which is normal for mineral materials". However, this argument is undermined by the fact that calcified tree roots very often form a 'column' stretching from ceiling to floor, yet a 'column' in a cave is a speleothem. Other secondary cave deposits such as flowstone, rimstone dam, cave pearls, dog tooth spar and many others similarly lack any '-ite' at the end of their name. Also, the term 'arborite' implies trees, from the Latin *arbor* 'tree'. The roots in caves may be from trees, but they may also be those of bushes, shrubs or even grass. In 1948, the word Arborite became a registered company name with manufacturing and distribution around the world, for a large product range of high-pressure urea formaldehyde laminates plus other products commonly referred to as arborite. So, for clarity the term 'rootsicle' is the best generic term for a root coated in a mineral deposit.

Conclusion

Given there are so many possible variables associated with roots that have been partly or completely coated by a secondary mineral, inside and outside caves, it would be logical to use the generic term "rootsicle", without defining the amount or thickness of mineral coating. If it occurs in a cave then it can be classed as a speleothem but if it occurs outside then it is not a speleothem. If a root grows through or around a pre-existing speleothem or above-ground secondary mineral deposit, it is not a rootsicle, unless it has a subsequent layer of secondary mineral deposited around the actual root.

Original definition (Smith 2010a)

ROOTSICLE. *n.* roots of trees or plants which grow into a cave cavity and become calcified. The roots and speleothem comprising the rootsicle.

Proposed new definition

ROOTSICLE. *n.* roots of trees or plants which have become partly or fully coated by a secondary mineral deposit, above or below ground. The roots (or remains of roots) and mineral deposit compris-

ing the rootsicle. If the root completely decays then the remaining secondary mineral skeleton or part thereof, is still considered to be a rootsicle.

If the rootsicle occurs inside a cave, then the mineral deposit is considered to be a speleothem, but if outside a cave it is not. If the growth of the speleothem (or above ground secondary mineral deposit) precedes the root, then it is not a rootsicle (refer Table 1).

Acknowledgements

Thanks to Cathi Humphrey-Hood and Rod OBrien for tracking down and scanning one of the reference articles held in the ASF Library. Also, thanks to Katerina Fulton for proofreading this paper.

References

- BAKER, A. 2007 Rock-me Cave. Saturday 25th November, 2006. *Newcaves Chronicles*, 27: 43.
- BASTIAN, L. 2014 Arborite: An alternative term for rootsicle. *The Western Caver*, 54: 4-6.
- BLANCHON, P. and SHAW, J. 1995 Reef drowning during the Last Deglaciation: Evidence for catastrophic sea-level rise and ice-sheet collapse. *Geology*, 23: 4-8.
- BUNNELL, D. The Virtual Cave: Rootsicles. <http://www.goodearthgraphics.com/virtcave/rootsicles/rootsicles.html> (accessed 27/12/2021).
- DAZA BRUNET, R. and BUSTILLO, M.A. 2014 Allophanic and ferric root-associated stalactites: Biomineralization induced by microbial activity (Galeria da Queimada lava tube, Terceira, Azores). *Geological Magazine*, 152(3): 504-520.
- DE WAELE, B. 2014 ... And what the other half did. *The Western Caver*, 54: 27-30.
- DU PREEZ, G.C., FORTI, P., JACOBS, G., JORDAAN, A. and TIEDT, L.R. 2015 Hairy stalagmites, a new biogenic root speleothem from Botswana. *International Journal of Speleology*, 44(1): 37-47. <http://dx.doi.org/10.5038/1827-806X.44.1.4>

- GILLIESON, D. 2004 Floral Resources [in] Gunn, J. (ed.) *Encyclopedia of caves and karst science*, Fitzroy Dearborn, New York, London. pp. 360-362.
- GONZÁLEZ-GONZÁLEZ, A.H., ROJAS-SANDOVAL, C., TERRAZAS, A., BENAVENTE, M., STINNESBECK, W., AVILES, O.J., DE LOS RÍOS, M. and ACEVEZ, E. 2008 The arrival of humans on the Yucatan Peninsula: Evidence from submerged caves in the State of Quintana Roo, Mexico. *Current Research in the Pleistocene. Special Report*, 25: 1-24.
- HILL, C. 1992 Mineralogy, sedimentology, and speleogenesis of Kartchner Caverns [in] BUECHER, R. (ed.) Final report: Environmental and geologic studies for Kartchner Caverns State Park. Arizona Conservation Projects, Tucson, 1: 271-301.
- HILL, C. 1999. Mineralogy of Kartchner Caverns, Arizona. *Journal of Cave and Karst Studies*, 61(2): 73-78.
- HILL, C. and FORTI, P. 1997 Related Forms: Rootsicles. *Cave Minerals of the World*. NSS, Huntsville, Alabama. Second edition, p. 224.
- JAMES, J.M., JENNINGS, J.N. and DYSON, H.J. 1982 Mineral decoration and weathering of the caves [in] DYSON, H.J., ELLIS, R. and JAMES, J.M. (eds.), *Wombeyan Caves*. Sydney Speleological Soc. Occasional Paper No. 8: 121-136.
- KLAPPA, C.F. 1980 Rhizoliths in terrestrial carbonates: classification, recognition, genesis and significance. *Sedimentology*, 27: 613-629.
- LAMONT, B. and LANGE, B. 1976 'Stalagmiform' roots in limestone caves. *New Phytologist*, 76: 353-360.
- MARAIS, J.C.E., IRISH, J. and MARTINI, J.E.J. 1996 Cave investigations in Namibia V: 1993 SWAKNO results. *Bulletin of the South African Speleological Association*, 36: 58-78.
- MLEJNEK, R. 2010 America, look for root stalagmites. *NSS News*, 68(4): 14-15.
- MLEJNEK, R., OUHRABKA, V. and RUZICKA, V. 2008 Poseidon – The pseudokarst systems in the sandstones of the Czech Republic: geomorphological structure and biogeographical importance. *Proceedings of the 10th International Symposium on Pseudokarst*, pp.75-86.
- MLEJNEK, R., OUHRABKA, V. and RUZICKA, V. 2009 Poseidon - a complex system of underground spaces in sandstone in the Czech Republic. *NSS News*, 67(8): 4-7.
- MOORE, G.W. 1952 Speleothems – a new cave term. *NSS News*, 10(6): 2.
- RUTLEDGE, J. 2008 Cave Descriptions and Maps [in] RUTLEDGE, J., SMITH, G.K., BRAINWOOD, M. and BAKER, A.C. (eds) *Timor Caves, Hunter Valley NSW*. N.&H.V.S.S., Broadmeadow, NSW. pp. 11-68.
- SHOPOV, Y.Y. 2004 Sediments: Biogenic [in] GUNN, J. (ed.) *Encyclopedia of Caves and Karst Science*. Fitzroy Dearborn, New York, London. pp. 636-637.
- SMITH, G.K. 1999 Glossary of Caving Terms – Common terminology and colloquial terms used throughout the speleological fraternity [in] *Proceedings of the Australian Speleological Federation 22nd Biennial Conference*, Yeppoon Qld. pp. 59-83.
- SMITH, G.K. 2001 Cliefden and Walli Caves 13th-16th April, 2001. *Newcaves Chronicles*, 17: 21-22.
- SMITH, G.K. 2003 West Australian caving - ASF Conference 2003, 'Underway'. *Newcaves Chronicles*, 20: 12-15.
- SMITH, G.K. 2007 Mt Gambier, SA: 26th Biennial Conference of the Australian Speleological Federation 6th to 12th Jan 2007 (Plus post field trips). *Newcaves Chronicles*, 28: 10-18.
- SMITH, G.K. 2009 Method of repairing the rootsicle in Wildmans Cave, Wombeyan 2- 3 May 2009. *Newcaves Chronicles*, 32: 27-29.

Rootsicles, roots & caves

- SMITH, G.K. 2010a Repairing the rootsicle in Wildmans Cave, Wombeyan, NSW. *Caves Australia*, 181: 18-20.
- SMITH, G.K. 2010b Glenrock Station, 12-14 June 2010. *Newcaves Chronicles*, 34: 32-25.
- SMITH, G.K. 2012 Timor and Crawney Pass caves, 6-8 April 2012. *Newcaves Chronicles*, 38: 14-17.
- SMITH, G.K. 2015 Jewboy Cave and Mt Sugarloaf 31 August 2014. *Newcaves Chronicles*, 43: 11-15.
- SMITH, G.K. 2018 Australia's largest cave systems in Judbarra-Gregory National Park, NT. *Caves Australia*, 206: 14-18.
- TABOROŠI, D., HIRAKAWA, K. and STAFFORD, R. 2004 Interactions of plant roots and speleothems. *Journal of Subterranean Biology*, 2: 43-51.
- TABOROŠI, D., HIRAKAWA, K. and SAWAGAKI, T. 2005 Carbonate precipitation along a microclimatic gradient in a Thailand cave - Continuum of calcareous tufa and speleothems. *Journal of Cave and Karst Studies*, 67(1): 69-87.
- TABOROŠI, D. 2006 Biologically influenced carbonate speleothems [in] HARMON R.S., WICKS, C.M., FORD, D.C. and WHITE, W.B. (eds) *Perspectives on Karst Geomorphology, Hydrology, and Geochemistry: A Tribute Volume to Derek C. Ford and William B. White*. Geological Society of America.
- WRAY, R.A.L. 1995 Solutional Landforms in Quartz Sandstones of the Sydney Basin. Thesis submitted to the School of Geosciences, University of Wollongong.
- WRAY, R.A.L. 1999 Opal and chalcedony speleothems on quartz sandstones in the Sydney region, southeastern Australia. *Australian Journal of Earth Sciences*, 46(4): 623-632.
- WINKELHOFER, R. 1975 Stalagmitenförmige Wurzelbindungen in Sandsteinhöhlen. *Der Höhlenforscher*, 7(2): 25-26.



**AUSTRALIAN
SPELEOLOGICAL
FEDERATION**